



Macro-level mental health system indicators and cross-national suicide rates

Johnny Andoh-Arthur & Samuel Adjorlolo

To cite this article: Johnny Andoh-Arthur & Samuel Adjorlolo (2021) Macro-level mental health system indicators and cross-national suicide rates, *Global Health Action*, 14:1, 1839999, DOI: [10.1080/16549716.2020.1839999](https://doi.org/10.1080/16549716.2020.1839999)

To link to this article: <https://doi.org/10.1080/16549716.2020.1839999>



© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 19 Jan 2021.



Submit your article to this journal [↗](#)



Article views: 2098



View related articles [↗](#)



View Crossmark data [↗](#)

Macro-level mental health system indicators and cross-national suicide rates

Johnny Andoh-Arthur ^c and Samuel Adjorlolo ^{a,b}

^aDepartment of Mental Health, School of Nursing and Midwifery, College of Health Sciences, University of Ghana, Accra, Ghana;

^bResearch and Grant Institute of Ghana, Accra, Ghana; ^cDepartment of Psychology, School of Social Sciences, University of Ghana, Accra, Ghana

ABSTRACT

Background: The relationship between macro-level mental health system indicators and population suicide rates is an area of contention in the literature, necessitating an analysis of current cross-national data to document any new trend in the relationship.

Objective: This study investigated whether mental health system indicators are associated with national suicide rates.

Method: Using an ecological study design and multivariate non-parametric robust regression models, data on suicide rates and mental health system indicators of 191 countries retrieved from WHO's 2017 Mental Health Atlas were compared.

Results: Findings revealed that the average suicide mortality rate was significantly higher in high-income countries, relative to low-income countries. High-income countries are significantly more likely to have high number of mental health professionals, mental health policies and legislation, independent mental health authority and suicide prevention programs. These mental health system indicators demonstrated significant and positive association with suicide, suggesting that countries scoring high on these factors have higher odds of being categorized as high suicide risk countries.

Conclusion: The findings have several implications for policy and practice, including the need to make existing mental health systems very responsive to suicide prevention.

ARTICLE HISTORY

Received 24 August 2020

Accepted 15 October 2020

RESPONSIBLE EDITOR

Jennifer Stewart Williams,
Umeå University, Sweden

KEYWORDS

Suicide; mental health;
mental health governance;
mental health professionals;
suicide prevention



Background

Suicide accounts for about 800,000 deaths annually at an estimated age standardized rate of 11.2 per 100,000 [1]. Across the globe, suicide is the 15th most common cause of death [1,2]. Suicide also reportedly accounted for 1.4% of premature deaths worldwide in 2015 [3]. A recent study revealed that the age standardized mortality rate for suicide has decreased by 32.7% worldwide between 1990 and 2016, however, total number of deaths from suicide within the same period increased by 6.7% globally over the 27-year-study period [4]. Suicide is notably a complex, multifactorial phenomenon that shows significant variations, particularly with respect to the causal factors and mechanisms, as well as prevention and management strategies [5].

Mental health problems such as mood, substance use, psychotic or personality disorders have been shown to play critical roles in suicide trajectory [6–11]. Often repeated are some psychological autopsy studies from High-Income Countries that for instance, suggest that 90%–95% of individuals who die of suicide had a diagnosable psychiatric disorders at the time of committing suicide [7,8,12]. Some authors have challenged this statistic by pointing either to methodological flaws that usually produces the statistic or to evidence suggesting no diagnosable psychiatric disorders in some

suicides in other contexts [13,14]. Indeed a systematic review from the Low-and Middle-Income countries (LMICs) has pointed to crucial role of non-psychiatric factors such as poverty in suicides [15]. It is possible that where mental health service is organized almost exclusively to focus on biomedical factors, it is most likely that less priority will be given to non-psychiatric factors in suicides [16]. Impliedly, cross-national variations in the role of psychiatric disorders and for that matter suicide rates generally could be partly attributed to variations in patterns, frequency, and meanings of suicide, as well as variations in the organization and provision of mental health service [6,16–18]. Among the factors that are likely to influence suicide rates across regions and countries are national income, mental health governance system, and resources for mental health, the so-called macro-level indicators of suicide because they require political commitment [19–21].

A number of ecological, country-level studies have examined the presumed link between suicide rate and mental health system indicators, such as availability of mental health services [5], professional density [2,22,23], mental health spending [24], mental health legislations and policies [21] and antidepressant sales [23]. A study conducted in Finland found that the prominence of outpatient services was associated

CONTACT Samuel Adjorlolo  sadjorlolo@ug.edu.gh  Department of Mental Health, School of Nursing and Midwifery, College of Health Sciences, University of Ghana, P. O. Box LG 43, Legon, Accra, Ghana

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

with lower suicide rate [17] whereas others have reported high suicide rates in countries with better mental health services such as higher numbers of psychiatric beds and availability of training in mental health for primary care professionals [5,18]. With respect to professional density and suicide rate, findings are mixed, with some studies reporting negative relationship [2,23,25,26], positive relationship [5,18,22], or no statistically significant relationship [23]. An Austrian study, for instance, found that a high number of mental health professionals was associated with lower rates of suicide, adjusting for per capita alcohol consumption and unemployment rates [23]. In contrast, cross-national data from 191 countries revealed that the number of psychiatrists was significantly positively associated with population suicide rates [5]. In yet another study from Austria, no statistically significant relationship was observed between psychiatrist density and suicide rate [23].

Analyses of cross-national data showed an increase in suicide rates and existence of mental health initiatives such as mental health policy, mental health program [18,21]. More importantly, while suicide rates reportedly differ across countries [6], the direction of the difference is an area of contention. More specifically, contrary to the finding that a vast majority, namely 78%, of suicides occur in the Low and Middle-Income Countries [3], others have found a high prevalence of suicide in high-income countries, relative to LMICs [27,28]. In their analysis of cross-national data [18], found that countries with high mental health expenditure tend to experience high suicide rate.

The inconsistent findings provide impetus for more studies to examine suicide and mental health systems. There is the pressing need to subject current cross-national data to empirical analysis to document any new trend in the relationship between mental health system and suicide rates across geographical regions [5,22]. Consequently, this study is designed to first and foremost investigate the distribution of suicide rates across countries based on their national income, and second adopt an ecological framework¹ to investigate the associations between suicide rate and mental health systems indicators (e.g. the number of professionals, availability of mental health policies)..

Methods

Data source

The data were extracted from the mental health Atlas 2017 published by the WHO for member states. The Atlas is designed to monitor and evaluate progress

made by member countries in achieving the WHO's Comprehensive Mental Health Action Plan by 2020. Among the core requirements of the Action Plan are for member countries to, for instance, strengthen mental health governance, render comprehensive and integrated mental health care, and institute mental health promotion and prevention strategies [29]. The Atlas, therefore, provides information on the state of mental health across nations by unearthing the existence or otherwise of mental health-specific indices pertinent to the Comprehensive Mental Health Action Plan. These include policies, plans and laws for mental health, human and financial resources; the type of mental health facilities, and availability of mental health promotion and prevention programs.

Data collection procedure

As noted previously, the Atlas is produced by the WHO with data from member countries. For the 2017 Atlas, data were collected using a structured questionnaire from 177 out of the 194 WHO member states, representing 97% of the world's population. The questionnaires were developed in consultation with member states and international experts in the area of mental health care measurement. In each member state, a focal person was identified to complete the questionnaire by extracting data from multiple sources, including from local team of experts, institutions such as Mental Health Authority/Commission and psychiatric and non-psychiatric hospitals located across different parts of the country. Once the focal person has completed the data-gathering process, the questionnaires were sent back to the WHO for processing and use. Where applicable, the focal persons were re-contacted for further information and clarifications relating to the questionnaire to ensure data quality. Detailed procedure for data collection has been described elsewhere [30].

Measures/instrument

Dependent variable

Suicide. The suicide mortality rate for the various countries were derived from the World Health Statistics data visualization dashboard, which is available at <http://apps.who.int/gho/data/>

Independent variables

The 2017 Atlas contain different macro-level data on mental health service delivery. Some variables (e.g. government's total expenditure on mental health, number of mental health hospitals, and number of

¹.....and second adopt an ecological framework to investigate the associations between suicide rate and mental health systems indicators (e.g., the number of professionals, availability of mental health policies). Ecological framework views problems i.e. suicide as the outcome of interaction among many factors at multiple levels – the individual, the relationship, the community, and the societal.

psychiatric units in a general hospital) were missing data at a scale that does not allow meaningful comparison. For example, in some instances, data are available for high-income countries but not of other income groups. Consequently, a decision was reached to exclude some data from the analysis. Variables with complete data across the various income groups were considered in this study, as stated below;

Income group. The 2017 Atlas categorized WHO member countries into four different income groups using the gross national income (GNI) developed by the World Bank in 2016. These are low income (GNI per capita of US\$ 1,025 or less), low-middle income (GNI per capita between US\$ 1,026 and US\$ 4,035), upper-middle income (GNI per capita between US\$ 4,036 and US\$ 12,475), and high income (GNI per capita of US\$ 12,476 or more).

Mental health system governance. The following variables were extracted to index mental health system governance: (1) Stand-alone policy or plan for mental health; (2) plan or strategy for child and/or adolescent mental health; (3) Stand-alone law for mental health; (4) existence of a dedicated authority or independent mental health body. Responses to these variables were present (Yes) or absent (No).

Resources for mental health. The variables extracted under resources for mental health were: (1) total number of mental health workers per 100,000 population; (2) number of psychiatrists; (3) number of mental health nurses; and (4) clinical psychologists per 100,000 population. Responses to these variables were continuous.

Suicide prevention programs. Data were collected on whether the countries have suicide prevention programs, with Yes or No as the response options.

Data analytic strategy

Data were summarized using descriptive statistics, notably frequencies, percentages, and bar chart. This was followed by inferential statistics using chi square (χ^2) and binomial logistic regression, with alpha level set at .05. The principal outcome variable, national suicide rates, and the resources for mental health variables did not follow Gaussian distribution. Consequently, we proceeded with the inferential statistical analysis by dichotomizing these variables [31]. First, they were numerically equalized and scaled along a common standard metric by converting them into z-scores. The standardized scores were subsequently recoded such that scores below and above the mean were designated as low (0) and high (1), respectively. The Yes/No responses to the

variables under the mental health system governance domain were also recoded as 1 and 0, respectively, for statistical purposes.

Next, χ^2 analysis was conducted to determine the association between the independent variables and suicides. Phi and Cramer's V coefficients were used to estimate the strength of the association, with coefficient values between .10 and .29, .30 to .49, and values at .50 and above interpreted as small, moderate and large effects. This was followed by a binomial logistic regression to determine the predictors of suicide categorization using odd ratios (OR) and adjusted odd ratio (AOR), where applicable. Income group and the variables under mental health system governance and resource for mental health domains were entered into the regression model independently. Only the variables demonstrating significant associations with suicide categorization based on χ^2 analysis were included in this analysis. Last, because national income is a major determinant of several indicators of mental health, including the propensity to increase the number of mental health professionals and enact mental health legislations [5], the effect of income group was controlled for to obtain the 'true effect' of mental health legislation and professionals on suicide categorization. All analyses were performed using SPSS version 24.

Results

National income and suicide

The average suicide mortality rate for 155 countries was 8.96 (Range: .50–31.90). Income group-based analyses revealed that low income ($n = 27$), low-middle income ($n = 42$), upper-middle income ($n = 46$) and high-income countries ($n = 40$) recorded 7.06 (range: 3.7–12.20), 7.61 (range: 1.90–22.40), 8.49 (range: 1.70–31.00) and 12.18 (.50–31.90) average suicide mortality rate, respectively. Income group and suicide rate are significantly correlated, $\chi^2(3) = 14.47$, $p < .01$, Phi & Cramer's V = .31. Countries recording at least 20 suicide rate per 100,000 include Belgium, Latvia, Ukraine, Suriname, South Korea, Guyana, Russian Federation, and Lithuania. In contrast, Antigua and Barbuda, Barbados, Grenada, Bahamas and Syrian Arab Republic were among the countries that reported less than two suicide rate per 100,000. From Table 1, national income and suicide rate evinced statistically significant relationship, $\chi^2(3) = 14.47$, $p < .05$. Further analyses showed that the odds of being designated as high suicide risk did not differ significantly between low income (the reference category) and low-middle and upper-middle income countries ($p > .05$). In contrast, high-income countries have higher odds of being classified as high suicide risk ($b = 1.67$, OR = 5.31).

Table 1. Relationship between national income, mental health system indicators and suicide.

Variables	Suicide			χ^2	P/C
	Low, <i>n</i> (%)	High, <i>n</i> (%)	Total, <i>n</i> (%)		
National Income				14.47***	.31
Low income	20(21.5)	7(11.3)	27(17.4)		
Low middle income	28(30.1)	14(22.6)	42(27.1)		
Upper middle income	31(33.3)	15(24.2)	46(29.7)		
High income	14(15.1)	26(41.9)	40(25.8)		
Total	93(60)	62(40)	155(100)		
Mental Health Professionals					
Total mental health professionals	73(90.1)	28(58.3)	101 (78.3)	17.92***	.37
Low	8(9.9)	20(41.7)	28(21.7)		
High	81(62.8)	48(37.2)	129(100)		
Total					
Psychiatrists	72(90)	21(42.9)	93(72.1)	33.57***	.51
Low	8(10)	28(57.1)	36(27.9)		
High	80(62)	49(38)	129(100)		
Total					
Mental health nurses	62(88.6)	19(50)	81(75)	19.54***	.43
Low	8(11.4)	19(50)	27(25)		
High	70(64.8)	38(35.2)	108(100)		
Total					
Clinical psychologists	65(94.2)	27(75)	92(87.6)	8.04**	.28
Low	4(5.8)	9(25)	13(12.4)		
High	69(65.7)	36(34.3)	105(100)		
Total					
Mental Health Governance System					
Mental policies or plans	8(8.6)	6(9.7)	14(9)	.05	.02
No	85(91.4)	56(90.3)	141(91)		
Yes	93(60)	62(40)	155(100)		
Total					
Mental health law	35(37.6)	21(33.9)	56(36.1)	.04	.04
No	58(62.4)	41(66.1)	99(63.9)		
Yes	93(60)	62(40)	155(100)		
Total					
Child/adolescent mental health strategy	61(65.6)	27(43.5)	88(56.8)	7.37**	.22
No	32(34.4)	35(56.5)	67(43.2)		
Yes	93(60)	62(40)	155(100)		
Total					
Mental health authority/commission	53(57)	22(35.5)	75(48.4)	6.89**	.21
No	40(43)	40(64.5)	80(51.6)		
Yes	93(60)	62(40)	155(100)		
Total					
Suicide prevention strategy				10.57**	.26
No	75(80.6)	35(56.5)	110(71)		
Yes	18(19.4)	27(43.5)	45(29)		
Total	93(60)	62(40)	155(100)		

P/C = Phi and Cramer's coefficient.

** = $p < .01$; *** = $p < .001$.

Mental health professionals and suicide

The average number of mental health professionals per 100,000 for 137 countries that supplied complete data was 37.61. When analyzed against income group, it was observed that the average number of mental health professionals differ across the income groups; low income (i.e. 1.58), lower-middle income (i.e. 7.33), upper-middle income (i.e. 43.61), and high income (i.e. 90.12) countries. Chi square analysis revealed that high-income countries are significantly likely to have more mental health professionals per 100,000, relative to low middle and low-income countries, $\chi^2(3) = 45.07$, $p < .001$, Phi & Cramer's $V = .58$. High-

income countries with the highest number of mental health professionals include Finland (i.e. 250.55), USA (i.e. 271.28), Brazil (i.e. 317.45), Costa Rica (i.e. 341.94), and Monaco (i.e. 405.41). The least number of mental health professionals was recorded in sub-Saharan Africa (SSA) countries such as Chad (i.e. 0.04), Guinea (i.e. 0.05), Central African Republic (0.15), Mali (i.e. 0.16), and Kenya (i.e. 0.19). Similar trend was observed when the analysis was focused on specific mental health professionals, namely psychiatrists, mental health nurses and clinical psychologists.

As can be seen in Table 2, the odds of being designated as high risk for suicide is significantly higher among countries with a high number of mental health professionals ($b = 1.88$, OR = 6.52), psychiatrists ($b = 2.49$, OR = 12.00), mental health nurses ($b = 2.05$, OR = 7.75) and clinical psychologists ($b = 1.69$, OR = 5.42). When the effect of national income was statistically controlled for, all the variables but clinical psychologists, retained their statistical significance ($p < .05$).

Mental health system governance and suicide

A total of 162 countries provided data on mental health system governance. A large proportion of the countries ($n = 147$, 90.7%) have instituted mental health policies or plans. More than half of the countries ($n = 105$, 64.8%) reported that they have stand-alone mental health laws, whereas 83 countries (51.2%) have a dedicated authority or independent mental health commission that oversees mental health activities in their respective countries. With respect to child/adolescent mental health, approximately 57% ($n = 92$) of the countries indicated they do not have plan or strategy for child/adolescent mental health. A statistically significant association was observed between income group and existence of plan or strategy for child/adolescent mental health, $\chi^2(3) = 17.23$, $p = .001$, Phi & Cramer's $V = .33$, dedicated mental health authority, $\chi^2(3) = 11.71$, $p = .008$, Phi & Cramer's $V = .27$, and stand-alone mental health law, $\chi^2(3) = 15.36$, $p = .002$, Phi & Cramer's $V = .31$, but not with mental health plans/policies $\chi^2(3) = 1.33$, $p = .722$, Phi & Cramer's $V = .09$.

Suicide rate correlated significantly with the availability of child/adolescent mental health strategies, $\chi^2(1) = 7.37$, $p < .01$, Phi & Cramer's $V = .22$, and the existence of independent mental health authority or commission, $\chi^2(1) = 6.89$, $p < .01$, Phi & Cramer's $V = .21$, but not with the availability of mental health plans/policies and mental health laws ($p > .05$). Further analyses reveal that, the odds of suicide rate is significantly higher in countries with child/adolescent mental health strategies ($b = .91$, OR = 2.47) and independent mental health authority ($b = .88$,

Table 2. Logistic regression of suicide on income group and mental health indicators.

Variables	<i>b</i>	95% CI Odd ratio (Unadjusted)	<i>B</i>	95% CI Adjusted Odd ratio
Income Group				
Low versus low-middle income	.36	1.43 (.49, 4.18)	-	-
Low versus upper-middle income	.32	1.38 (.48, 3.99)	-	-
Low versus high income	1.67**	5.31 (1.81, 3.60)	-	-
Mental Health Personal (per 100,000)				
Total mental health professionals	1.88***	6.52 (2.56, 5.49)	1.50**	4.49 (1.47, 4.73)
Psychiatrists	2.49***	12.00 (4.76, 6.23)	2.71***	15.04 (4.23, 7.41)
Nurses	2.05***	7.75 (2.93, 4.50)	1.31**	6.10 (1.93, 3.29)
Psychologists	1.69**	5.42 (1.54, 3.10)	1.25	3.50 (.90, 3.69)
Mental health professionals				
Child/adolescent strategy	.91**	2.47 (1.28, 4.78)	.68	1.98 (.97, 4.05)
Authority/Commission	.88**	2.41 (1.24, 4.67)	.73*	2.07 (1.01, 4.21)
Suicide prevention programs				
	1.17**	3.21(1.57, 6.60)	1.06**	2.87(1.31, 5.31)

** = $p < .01$; *** = $p < .001$.

OR = 2.41). When the effect of national income was controlled for, only the existence of independent mental health authority significantly increased the odds of being labelled as high risk for suicide ($b = .73$ AOR = 2.07).

Suicide prevention programs and suicide

Of the 162 countries providing data on suicide, majority ($n = 116$, 71.6%) indicated they do not have stand-alone, government initiated national suicide prevention programs/plan. Income group correlated significantly with the existence of suicide prevention strategies, $\chi^2(3) = 15.34$, $p < .01$, Phi & Cramer's $V = .31$, with high-income group more likely to have national suicide prevention strategies. Indeed, when the analysis was disaggregated by income group, it was observed that only four low income ($n = 27$; i.e. Afghanistan, Mozambique, Chad and Uganda), five low-middle income ($n = 43$; e.g. Philippines, Vanuatu, Timor-Lesta, Bhutan and Nicaragua), 18 upper middle income ($n = 49$; e.g. Iran, Panama, Malaysia, Ecuador and Turkey) and 19 high-income countries ($n = 43$; e.g. Monaco, Israel, Italy and Spain) have national suicide prevention plans/strategies.

As shown in Table 1, availability of national suicide prevention plans was significantly associated with suicide rate, $\chi^2(1) = 10.57$, $p < .01$, Phi & Cramer's $V = .26$. Further analysis revealed that countries with suicide prevention strategies are significantly more likely to have higher odds of being categorized as high suicide risk country ($b = 1.17$, OR = 3.21), even after controlling for national income groupings ($b = 1.06$, AOR = 2.87).

Discussion

The study primarily investigated the macro-level factors influencing suicide mortality rates across countries using data from the WHO's 2017 Mental Health Atlas.

Suicide rates in low and middle income and high-income countries

While suicide remains one of the global challenges confronting nations, the study found that the average suicide mortality rate was significantly higher in high-income countries, relative to low-income countries. This finding, which is largely consistent with previous studies [18,27,28], contradicts the widely held view that LMICs tend to experience the greatest proportion of suicides [3]. The supposedly low prevalence of suicide in LMIC could be accounted for by several factors, including the relatively high social support, religious commitment and/or involvement, and better family cohesion, which are potential protective factors against suicide [32,33].

The foregoing notwithstanding, there is also the possibility that real-suicide figures are obscured in LMIC partly due to the fundamental problem of inaccurate or lack of data pertaining to suicide. Indeed, suicide rates depend not only on the efficiency of civil registration systems, which are generally poor in LMIC, but also on the reporting of deaths which is in turn heavily influenced by the social, cultural and legal consequences of suicide [34]. Despite the ongoing campaign and advocacy to decriminalize suicide across countries, suicide continues to exist as a legal term that is often accompanied by legally prescribed punitive measures in several sub-Saharan African countries, including Ghana [35], Nigeria, Botswana, Gambia, Kenya, Malawi, Tanzania, Zambia and Ugandan [36]. In some LMICs such as Ghana, suicide is viewed as a taboo and unnatural death. Given this orientation, individuals expressing suicidal tendencies or persons deceased by suicides as well as their families are socially sanctioned. This observation incentivize families and communities to conceal or misreport suicidal behaviors so as to protect and preserve the sanctity of the family name [37,38]. Under-reporting of suicide is therefore highly prevalent in LMICs [28,39] owing to the prevailing social, cultural and legal proscriptions against suicides [5].

In contrast, the reported high prevalence of suicide in high-income countries could be due to an artifact of more efficient death registration systems and their case finding effects [34]. Likewise, the implementation of suicide prevention programs and the associated awareness and hypervigilance could culminate into case-finding effect where there is the general tendency to label deaths as suicides [21].

The foregoing has enormous implication on the use of population suicide rate as a proxy indicator of the effectiveness of a country's mental health services. That is, although LMIC reportedly have low suicide rates, it will be problematic to attribute this to better mental health system in these countries, as discussed previously. Indeed, LMICs are characterized by low number of mental health professionals, relative to high-income countries; a development that has been attributed to low investment in training mental health professionals and the relocation of mental health professionals to high-income countries for better conditions of service. For instance, on the latter point, an earlier study reported more Ghanaian psychiatrists working in the USA than those working in Ghana [40].

Correlates of suicides rate

The study also found that high-income countries are significantly more likely to have child and adolescent mental health strategies, dedicated mental health authority and suicide prevention programs, relative to LMIC. More importantly, given the robust link between mental health problems and tendency for suicide particularly in the high-income countries [9,11], it is plausible to reason that programs and initiatives designed to promote and restore mental health functioning will contribute to a reduction in suicide rate. In contrast, however, the study found that macro-level mental health indicators examined in this study (e.g. number of mental health professionals, dedicated mental health authority and national suicide prevention programs) are associated with an increase in suicide rate, even after controlling for national income. This is largely consistent with previous studies [5,21]. Although the finding is somewhat counterintuitive and so could discourage further investment in mental health, on the other hand, it presents another opportunity for policy makers and relevant stakeholders to reexamine the various national mental health initiatives within the ambit of suicide prevention [21]. More specifically, the study highlighted several possibilities, including the view that the aforementioned national mental health initiatives may not be designed with a focus on suicide prevention, notwithstanding that they are intended to promote general mental health and well-being. Moreover, the over-medicalization of suicide,

predominantly in high-income countries implies that greater attention is paid to urgency-driven curative medical solutions, thereby belittling the importance of associated sociocultural and economic factors [5,41]. Indeed, the observation that suicide can occur in the absence of psychological and mental health problems [42] implies that programs mainly designed to promote and improve mental health in general may have contributed little to reducing the prevalence of suicide rates. Some major risk factors that have been found to have exhibited direct and indirect relationship with suicide include unemployment, alcohol consumption and substance misuse in general, social inequalities, loss of social cohesion, and financial difficulties emanating from high indebtedness and bankruptcy [43–45]. An earlier study, for instance, found that the relationship between unemployment and suicide was statistically significant even after controlling for mental illness [46]. The foregoing presupposes that appropriate macroeconomic and social welfare policies that ensure and promote basic human rights, social security, gender equality and equitable development may contribute to reducing population suicide rates [5]. Perhaps, it is against this background that others have renewed calls for situating suicide research and prevention within social and cultural contexts [47]. The purpose is to deepen the understanding of suicides in accordance with postulation of the stress-diathesis model suggesting that suicide risk is multi-factorial [48] and should therefore not be reduced to only psychiatric illness [6]. Furthermore, although specific suicide prevention programs may exist particularly in high-income countries, there is the possibility they have not been implemented [49], or even if they are implemented, they could be riddled with implementation issues that could thwart or reduce their effectiveness.

Limitations

The findings of the study should be reviewed in light of the following limitations. First, all ecological studies have the potential limitation of ecological fallacy, which can occur as an association observed between the study variables on the aggregate level will not necessarily represent the association existing at the individual level. Causal associations cannot be assumed since this study analyzed data collected using a cross-sectional study design. The study could also be confounded by potential variables such as cultural and religious differences between countries which were not captured in the WHO's 2017 Mental Health Atlas. There is also the possibility of under or over-reporting of data which may occur for several reasons, including an attempt to present a good picture of mental health situations. The application of advanced data mining techniques may uncover patterns that might have

been missed with the data analytic strategy employed in this study. Also, because the quality of training and focus of psychiatrists might differ across countries [22], the lack of data on these critical variables could affect the findings of the study. Last, the findings reported in this study could have been different with data from all WHO member countries. Notwithstanding the limitations, the findings reported in this study are largely consistent with previous ecological studies.

Conclusion

This ecological study examined the association between population suicide rates and key mental health indicators based on the most recent WHO Mental Health Atlas which reported suicide rates from 191 countries. Findings generally revealed significant and positive relationship between macro-level mental health indicators and suicides. Although the findings appear somewhat counterintuitive in the light of the robust link between suicide and mental illnesses in many countries, nonetheless, they prompt action towards improving suicide reporting and recording systems across the globe generally and mapping of locally relevant factors contributing to suicides as a guide for formulating and implementing context sensitive prevention and intervention measures. .

Noteworthy is the lower rate of suicides in individual countries within the LMICs, which is as revealing as it is intriguing. Suicide and suicide prevention within most LMIC countries are not national priorities [50]. Apart from the social, cultural and legal barriers that hinder accurate reporting and recording of suicides, it is possible that the burden of disease attributable to communicable and infectious diseases in LMICs tended to sway concerted national health responses away from non-communicable and relatively ‘rare’ health problems such as suicides. Where modest attention was given to the latter, it was largely focused on cardiovascular diseases, diabetes, cancer and chronic pulmonary diseases [51,52]. However, given that some countries in the LMICs have or are undergoing ‘epidemiological transition’ whereby the relative importance of infectious diseases is gradually becoming less than that of chronic diseases [53], there is a need for national governments within the LMICs to commit more resources to non-infectious causes of mortality and morbidity such as suicides. Such efforts will be in keeping with the target 3.4 of the Sustainable Development Goals, which is to reduce by one-third premature mortality from non-communicable diseases by the year 2030 [54]. In the light of the above, we suggest the need to increase qualitative and quantitative research within individual countries that will help deepen understanding of the meaning of suicide and identifiable risk factors for people and how and why

common risk factors contribute to suicides in some people but not in others.

Acknowled

We are grateful to Mr. Godfred Danso for extracting the data from the WHO Mental Health Atlas

Author contributions

SA conceived the idea and extracted data; JAA drafted the manuscript, SA performed the analysis and wrote result, JAA wrote the discussion and conclusion; SA and JAA proofread the manuscript.

Disclosure statement

No conflict of interest was reported by the authors

Ethics and consent

The study used data generated by the WHO

Funding information

There is no external funding to declare

Paper context

Despite the robust association between mental health status and risk for suicide at the individual level, ecological studies exploring mental health systems and suicide rates are not only scarce but have produced inconsistent results. The study found that (high income) countries with proper mental health system and governance are more likely to record high suicide rates. There is the need to make existing mental health systems very responsive to suicide prevention.

ORCID

Johnny Andoh-Arthur  <http://orcid.org/0000-0002-7036-1835>

Samuel Adjorlolo  <http://orcid.org/0000-0001-9308-6031>

References

- [1] WHO. Preventing suicide: global imperative. Geneva; 2014. [cited 2020 Oct 8]. Available from: https://www.who.int/mental_health/suicide-prevention/world_report_2014/en/
- [2] Kawaguchi H, Koike S. Association between the density of physicians and suicide rates in Japan: nationwide ecological study using a spatial Bayesian model. *PLoS One*. 2016;11:e0148288.
- [3] Bachmann S. Epidemiology of suicide and the psychiatric perspective. *Int J Environ Res Public Health*. 2018;15:1425.
- [4] Naghavi M. Global, regional, and national burden of suicide mortality 1990 to 2016: systematic analysis for the Global Burden of Disease Study 2016. *BMJ*. 2019;364. DOI:10.1136/bmj.194

- [5] Rajkumar A, Brinda E, Duba A, et al. National suicide rates and mental health system indicators: an ecological study of 191 countries. *Int J Law Psychiatry*. 2013;36:339–342.
- [6] Brådvik L. Suicide risk and mental disorders. *Int J Environ Res Public Health*. 2018;15:2028.
- [7] Bertolote JM, Fleischmann A. Suicide and psychiatric diagnosis: a worldwide perspective. *World Psychiatry*. 2002;1:181.
- [8] Arsenault-Lapierre G, Kim C, Turecki G. Psychiatric diagnoses in 3275 suicides: a meta-analysis. *BMC Psychiatry*. 2004;4:37.
- [9] Harris EC, Barraclough B. Suicide as an outcome for mental disorders. A meta-analysis. *Br J Psychiatry*. 1997;170:205–228.
- [10] Andoh-Arthur J, Hjelmeland H, Osafo J, et al. Substance use and suicide among men in Ghana: a qualitative study. *Curr Psychol*. 2020;1–13. DOI:10.1007/s12144-020-00644-0
- [11] Furqan Z, Sinyor M, Schaffer A, et al. “I can’t crack the code”: what suicide notes teach us about experiences with mental illness and mental health care. *Can J Psychiatry*. 2019;64:98–106.
- [12] Cavanagh JT, Carson AJ, Sharpe M, et al. Psychological autopsy studies of suicide: a systematic review. *Psychol Med*. 2003;33:395–405.
- [13] Hjelmeland H, Dieserud G, Dyregrov K, et al. Psychological autopsy studies as diagnostic tools: are they methodologically flawed? *Death Stud*. 2012;36:605–626.
- [14] Shahtahmasebi S. Suicide prevention: politics or conspiracy. *Dyn Hum Behav (DHH)*. 2016;3.
- [15] Iemmi V, Bantjes J, Coast E, et al. Suicide and poverty in low-income and middle-income countries: a systematic review. *Lancet Psychiatry*. 2016;3:774–783.
- [16] Boldt M. The meaning of suicide: implications for research. *Crisis*. 1988;9:93–108.
- [17] Pirkola S, Sund R, Sailas E, et al. Community mental-health services and suicide rate in Finland: a nationwide small-area analysis. *Lancet*. 2009;373:147–153.
- [18] Shah A, Bhandarkar R, Bhatia G. The relationship between general population suicide rates and mental health funding, service provision and national policy: a cross-national study. *Int J Social Psychiatry*. 2010;56:448–453.
- [19] Homan C, Johar R, Liu T, et al., editors. Toward macro-insights for suicide prevention: analyzing fine-grained distress at scale. *Proceedings of the Workshop on Computational Linguistics and Clinical Psychology: From Linguistic Signal to Clinical Reality*; 2014. [cited 2020 Oct 8]. Available from: <https://www.aclweb.org/anthology/W14-3213.pdf>
- [20] Lee J. Municipal-level determinants of suicide rates in South Korea: exploring the role of social capital and local government policies. *J Asian Public Policy*. 2019;1–18. DOI:10.1080/17516234.2019.1565336
- [21] Burgess P, Pirkis J, Jolley D, et al. Do nations’ mental health policies, programs and legislation influence their suicide rates? An ecological study of 100 countries. *Aust N Z J Psychiatry*. 2004;38:933–939.
- [22] Sher L. Are suicide rates related to the psychiatrist density? A cross-national study. *Front Public Health*. 2016;3. DOI:10.3389/fpubh.2015.00280
- [23] Kapusta ND, Niederkroenthaler T, Etzersdorfer E, et al. Influence of psychotherapist density and antidepressant sales on suicide rates. *Acta Psychiatr Scand*. 2009;119:236–242.
- [24] Ross JM, Yakovlev PA, Carson F. Does state spending on mental health lower suicide rates? *J Socio Econ*. 2012;41:408–417.
- [25] Korosec Jagodic H, Rokavec T, Agius M, et al. Availability of mental health service providers and suicide rates in Slovenia: a nationwide ecological study. *Croat Med J*. 2013;54:444–452.
- [26] Tondo L, Albert MJ, Baldessarini RJ. Suicide rates in relation to health care access in the USA: an ecological study. *J Clin Psychiatry*. 2006;67:517–523.
- [27] Khazaei S, Armanmehr V, Nematollahi S, et al. Suicide rate in relation to the Human Development Index and other health related factors: a global ecological study from 91 countries. *J Epidemiol Glob Health*. 2017;7:131–134.
- [28] Blasco-Fontecilla H, Artieda-Urrutia P, de Leon J. A proposal for using the ratio of attempted to completed suicides across several countries worldwide. *Epidemiol Psychiatr Sci*. 2018;1–5. DOI:10.1017/S2045796018000628
- [29] WHO. Mental health action plan 2013–2020. Geneva; 2013. [Cited 2020 Oct 8]. Available from: https://apps.who.int/iris/bitstream/handle/10665/89966/9789241506021_eng.pdf?sequence=1
- [30] WHO. Mental Health Atlas. Geneva; 2017. [Cited 2020 Oct 8]. Available from: <https://apps.who.int/iris/bitstream/handle/10665/272735/9789241514019-eng.pdf?ua=1>
- [31] Lumley T, Diehr P, Scott Emerson A, et al. The Importance of the normality assumption in large public health data sets. *Annu Rev Public Health*. 2002;23:151–169.
- [32] Rushing NC, Corsentino E, Hames JL, et al. The relationship of religious involvement indicators and social support to current and past suicidality among depressed older adults. *Aging Ment Health*. 2013;17:366–374.
- [33] Šedivy NZ, Podlogar T, Kerr DC, et al. Community social support as a protective factor against suicide: A gender-specific ecological study of 75 regions of 23 European countries. *Health Place*. 2017;48:40–46.
- [34] Joseph A, Abraham S, Muliylil J, et al. Evaluation of suicide rates in rural India using verbal autopsies, 1994–9. *BMJ*. 2003;326:1121–1122.
- [35] Adinkrah M. Criminal prosecution of suicide attempt survivors in Ghana. *Int J Offender Ther Comp Criminol*. 2013;57:1477–1497.
- [36] Adinkrah M. Anti-suicide laws in Nine African countries: criminalization, prosecution and penalization. *Afr J Criminol Justice Stud*. 2016;9:279–291.
- [37] Andoh-Arthur J, Hjelmeland H, Osafo J, et al. Walking a tightrope: reflections on police gatekeeping roles in suicide research in Ghana. *Int J Soc Res Methodol*. 2018;21:289–301.
- [38] Osafo J, Hjelmeland H, Akotia C, et al. Social injury: an interpretative phenomenological analysis of the attitudes towards suicide of lay persons in Ghana. *Int J Qual Stud Health Well-being*. 2011;6:8708.
- [39] Reynders A, Scheerder G, Van Audenhove C. The reliability of suicide rates: an analysis of railway suicides from two sources in fifteen European countries. *J Affect Disord*. 2011;131:120–127.

- [40] Doku VCK, Mallett MR. Collaborating with developing countries in psychiatric research. *Br J Psychiatry*. 2018;182:188–189.
- [41] Pridmore S. Medicalisation of suicide. *Malays J Med Sci*. 2011;18:78.
- [42] Milner A, Sveticic J, De Leo D. Suicide in the absence of mental disorder? A review of psychological autopsy studies across countries. *Int J Social Psychiatry*. 2013;59:545–554.
- [43] Andoh-Arthur J, Knizek BL, Osafo J, et al. Suicide among men in Ghana: the burden of masculinity. *Death Stud*. 2018;42:658–666.
- [44] Asante KO, Kugbey N, Osafo J, et al. The prevalence and correlates of suicidal behaviours (ideation, plan and attempt) among adolescents in senior high schools in Ghana. *SSM Popul Health*. 2017;3:427–434.
- [45] Haw C, Hawton K, Gunnell D, et al. Economic recession and suicidal behaviour: possible mechanisms and ameliorating factors. *Int J Social Psychiatry*. 2015;61:73–81.
- [46] Goldman-Mellor SJ, Saxton KB, Catalano RC. Economic contraction and mental health: a review of the evidence, 1990–2009. *Int J Ment Health Syst*. 2010;39:6–31.
- [47] Hjelmeland H, Loa Knizek B. The emperor's new clothes? A critical look at the interpersonal theory of suicide. *Death Stud*. 2018;1–11. DOI:10.1080/07481187.2018.1527796
- [48] Mann JJ, Waternaux C, Haas GL, et al. Toward a clinical model of suicidal behavior in psychiatric patients. *Am J Psychiatry*. 1999;156:181–189.
- [49] Jacob K, Sharan P, Mirza I, et al. Mental health systems in countries: where are we now? *Lancet*. 2007;370:1061–1077.
- [50] Jacob K. Suicide in india: part perceptions, partial insights, and inadequate solutions. *Natl Med J India*. 2017;30:155–158.
- [51] Vos T, Barber RM, Bell B, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;386:743–800.
- [52] Boutayeb A. The double burden of communicable and non-communicable diseases in developing countries. *Trans R Soc Trop Med Hyg*. 2006;100:191–199.
- [53] Phillips M. Suicide prevention in developing countries: where should we start? *World Psychiatry*. 2004;3:156.
- [54] UNDP. Sustainable development goals. n.d. [cited 2020 Apr 03]. Available from: <https://www.undp.org/content/undp/en/home/sustainable-development-goals.html>